

Figure 1. Map showing locations of samples collected by the USGS and used to determine distribution of sea-floor sediments.

Alaskan coastline, alaska.cst (USGS). <http://walrus.wr.usgs.gov/infobank/bear/poly/ak/alaska.cst>

Bathymetry modified from Etopo2 database (Smith and Sandwell, 1997).

Base grid plotted with Generic Mapping Tools, Version 3.0 (Wessel and Smith, 1995).

Mercator projection


NOTE: THIS MAP IS NOT INTENDED FOR NAVIGATION

MAP OF DISTRIBUTION OF BOTTOM SEDIMENTS ON THE CONTINENTAL SHELF, GULF OF ALASKA

By

Kevin R. Evans, Paul R. Carlson, Monty A. Hampton, Michael S. Marlow, and Peter W. Barnes

2000



Distribution of bottom sediments digitized by K.R. Evans, 1997, from Carlson and others (1977) and (1986).

The authors gratefully acknowledge the assistance given by Carolyn Degnan for geographic and bathymetric basemaps, and by Christina Guttmacher for final editing changes in preparation for publication.

This map was printed on an electronic plotter directly from digital files. Dimensional calibration may vary between electronic plotters and paper, and may not be the same for the same plotter, and paper may change size due to atmospheric conditions; therefore, scale and proportions may not be exact on plots of the same size.

For sale by U.S. Geological Survey, Map Distribution, Box 25286, Federal Center, Denver, CO 80225

Available on World Wide Web at <http://geopubs.wr.usgs.gov/map-mf2335>