


CORRELATION OF MAP UNITS



DESCRIPTION OF MAP UNITS


VERY YOUNG SURFICIAL DEPOSITS—Sediment recently transported and deposited in channels and washes... QUATERNARY: Qf, Qy, Qm, Qs, Qv, Qx, Qz, Tm, Trg, Ktt, Kgt, Kgc, Kkv, Kk, Kp, Kq, Kx, Kz, Kc, Kd, Ke, Kf, Kg, Kh, Ki, Kj, Kl, Km, Kn, Ko, Kp, Kq, Kr, Ks, Kt, Ku, Kv, Kw, Kx, Ky, Kz. TERTIARY: Tm, Trg, Ktt, Kgt, Kgc, Kkv, Kk, Kp, Kq, Kx, Kz. CRETACEOUS: Qm, Qy, Qz, Tm, Trg, Ktt, Kgt, Kgc, Kkv, Kk, Kp, Kq, Kx, Kz. MESOZOIC: Qm, Qy, Qz, Tm, Trg, Ktt, Kgt, Kgc, Kkv, Kk, Kp, Kq, Kx, Kz.

GEOLOGIC SUMMARY

The Steele Peak 7.5' quadrangle is located in the northern part of the Peninsular Ranges Province within the central part of the Perris block, a relatively stable, rectangular in plan area located between the Elmore and San Jacinto fault zones. The quadrangle is underlain by Cretaceous and older basement rocks. Cretaceous plutonic rocks are part of the composite Peninsular Ranges batholith. A wide variety of mafic to intermediate composition granitic rocks occur in the quadrangle, and are mainly of anatectic composition, but range from monzogranite to gabbro. Most rock units are fairly to intensely foliated, compositionally heterogeneous, and contain varying amounts of meso- and melanocratic inclusions. Some rocks are composed almost wholly of inclusion material and some are migmatitic. Included within these granitic rocks are some that are not shown on the geologic map (Paleozoic?) schist of upper amphibolite metamorphic grade. Metamorphic rocks of primarily Mesozoic age occur in a discontinuous belt extending from the southeast to the northeast corner of the quadrangle. Most of these rocks are well foliated biotite-bearing schist. Near the southern edge of the quadrangle phyllite rocks dominate. Northwestward, metamorphism increases from greenschist or sub-greenschist grade near the south edge of the quadrangle to sillimanite-bearing schist of upper amphibolite grade in the vicinity of Cajalco Road. Biotite-hornblende tonalite of the relatively large Val Verde pluton dominates the northeastern half of the quadrangle. It most places this tonalite has a northeast-southwest-trending well developed planar fabric produced by oriented biotite and hornblende. Schlieren and massive cuts of mafic tonalite locally occur. Discordant biotite-shaped mafic inclusions are widespread and are oriented in the plane defined by the biotite and hornblende. This planar fabric typically dips moderately to the northeast, but locally shallowly to a horizontal to subhorizontal planar fabric, or folds to an isoclinal fabric. West of the Val Verde pluton a number of plutons having fabrics ranging from massive isotropic to foliated. Compositions of these plutons range from monzogranite to pyroxene gabbro. Most of these granitic rocks fall within the composition range from monzogranite to tonalite, and are part of the composite Gavilan ring complex. Hypersthene is a characteristic mineral of most of the rocks of this complex, which include black hypersthene-bearing monzogranite that has been quarried in a source of black granite building stone. Several inactive gold mines, e.g., Goodhope, Gavilan, and Santa Rosa mines that constituted the Pinacate mining district, are located in the Gavilan ring complex. In the center of the Gavilan ring complex is the near-circular Arroyo del Toro pluton, a massive-textured granodiorite essentially devoid of inclusions. Only the northern half of this pluton is located in the quadrangle. Some rock of this pluton was quarried for building stone. The southwestern corner of the quadrangle is underlain by the Elmore and the Perris faults, and is characterized by highly deformed rocks in the upper part of the batholithic magmatism. Most of these volcanic rocks range in composition from rhyolite to andesite with latitic composition rocks predominating. In the northeastern part of the quadrangle is the proximal parts of a Pleistocene alluvial fan complex.

REFERENCES

Anderson, C.L., 1991. Zircon uranium-lead isotopic ages of the Santiago Peak volcanic and spatially related plutons of the Peninsular Ranges batholith, southern California, U.S. thesis, San Diego, California, San Diego State University, p. 111. Dudley, P.H., 1955. Geology of a portion of the Perris block, southern California. California Jour. of Mines and Geology, v. 31, no. 4, p. 487-506. Larsen, E.S., 1948. Batholith and associated rocks of Corona, Elmore, and San Luis Rey quadrangles, southern California. Geol. Soc. of America Mem. 29, 182 p. Morton, D.M., 1999. Preliminary digital geologic map of the Santa Ana 30' X 60' quadrangle, southern California. U.S. Geological Survey Open-File Report 99-172, 61 p., scale, 1:100,000. Osborn, J.F., 1939. Structural geology of the Val Verde Tonalite, southern California. Geol. Soc. America Bull., v. 50, p. 921-950. Rogers, T.H., 1965. Santa Ana sheet. California Division of Mines and Geology Geologic Map of California, scale, 1:240,000. Sampson, R.J., 1935. Mineral resources of a portion of the Perris block, Riverside County, California. California Journal of Mines and Geology, v. 31, p. 507-521. Streckenbach, A.L., 1973. Plutonic rocks—Classification and nomenclature recommended by the IUGA Subcommission on Systematics of Igneous Rocks. Geotitles, vol. 18, p. 26-30. Woodfield, A.O., McCallith, T.H., and Schoellhamer, J.E., 1973. Paleogeographic significance of mineral boulders in middle-Tertiary strata, Santa Ana Mountains, California. Geol. Soc. America Bull., v. 83, no. 11, p. 2435-2436. Yerkes, R.F., 1972. Geology and oil resources of the western Puente Hills area, southern California. U.S. Geol. Surv. Prof. Paper 420-C, 63 p.



GEOLOGIC MAP OF THE STEELE PEAK 7.5' QUADRANGLE, RIVERSIDE COUNTY, CALIFORNIA

Version 1.0
By
Douglas M. Morton
Digital preparation by
Rachel M. Alvarez and Van M. Diep

U.S. Geological Survey
Department of Earth Sciences
University of California
Riverside, CA 92521

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, firm, or product names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.
This map was printed on an electronic plotter directly from digital files. Dimensional calibration may vary between electronic plotters and between X and Y directions on the same plotter, and paper may change size due to atmospheric conditions; therefore, scale and proportions may not be true on plots of this map.
Digital files available on World Wide Web at http://geopubs.usgs.gov