



# Logs and Data from Trenches Across the Hayward Fault at Tyson's Lagoon (Tule Pond), Fremont, Alameda County, California, 2001- 2003

By James J. Lienkaemper<sup>1</sup>, Patrick L. Williams<sup>2</sup>, Timothy E. Dawson<sup>1</sup>, Stephen F. Personius<sup>3</sup>, Gordon G. Seitz<sup>4</sup>, Samuel J. Heller<sup>4</sup>, and David P. Schwartz<sup>1</sup>

Open-File Report 03-488

2003

This report is preliminary and has not been reviewed for conformity with U.S. Geological Survey editorial standards or with the North American Stratigraphic Code. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

**U.S. DEPARTMENT OF THE INTERIOR  
U.S. GEOLOGICAL SURVEY**

<sup>1</sup>U.S. Geological Survey, Menlo Park, CA 94025

<sup>2</sup>P. Williams & Assoc., P.O. Box 1492 West Tisbury MA 02575

<sup>3</sup>U.S. Geological Survey, Denver, CO 80225

<sup>4</sup>Lawrence Livermore National Laboratory, Livermore, CA 94551

## INTRODUCTION

The purpose of this publication is to make available detailed trench logs (sheets 1 through 8), radiocarbon dates (table 1) and brief stratigraphic unit descriptions (Appendix 1) obtained as a result of an intensive subsurface investigation of the Hayward Fault at Tyson's Lagoon (Tule Pond) from 2001 through 2003 (figs. 1, 2 and 3 on sheet 2). Preliminary findings of this investigation based on fieldwork done in 2000 have been published as Lienkaemper and others (data archive, 2002a; report, 2002b). The Hayward Fault is recognized to be among the most hazardous in the United States (Working Group on California Earthquake Probabilities, 1999). This document makes available geologic evidence for historical and prehistoric surface-rupturing earthquakes preserved at the site. Establishing a chronology of prehistoric or paleoearthquakes is of immediate use in resolving the level of hazard posed by the Hayward Fault for producing large earthquakes in the future. A second formal report on our conclusions based on these data is in preparation. The investigation at Tyson's Lagoon is ongoing, so these products should not be considered final. Lienkaemper, Williams, Dawson and Personius interpreted the geology and logged the trenches. Seitz and Heller performed analyses on the radiocarbon samples. Schwartz led the critical-review field team.

Previous trenching work was done at Tyson's Lagoon (figs. 2, 3 on sheet 2). Lienkaemper (1992) references the location of most of those trenches. The earlier trenching was generally for the evaluation of local fault-rupture hazard, except for the study of Williams (1993), which was a paleoearthquake investigation. An unpublished study by J.N. Alt in 1998 (shown on our site map as trenches 98A and 98B, fig. 3, on sheet 2), also sought evidence of paleoearthquakes. Alt's study and one by Woodward-Clyde and Associates (1970; trenches 70A to 70G, fig. 3) were located south of Walnut Avenue in one of the few areas that still remain undisturbed and were, thus, useful in planning our work in 2000. Unpublished field investigations at this site by P. L. Williams in 1992 have been incorporated in this report. His trench W2 is presented herein as sheet 8 (see fig. 3 sheet 2 for location). William's 1992 trench W1 was re-excavated by us in 2002 and is shown as trench 02E (sheet 4).

## METHODOLOGY

### TRENCHING

As seen on the map, Tyson's Lagoon (fig. 2 on sheet 2) is a classic example of a sag pond (*e.g.*, Wallace, 1990), a pull-apart structure caused by a right step in a right-lateral fault. We excavated one trench in 2001, 01A (sheet 1); five trenches in 2002, 02A (sheet 2), 02B, C, D (sheet 3) and 02E (sheet 4); and two trenches in 2003, 03A (sheet 5) and 03B (sheet 6). The site location maps showing the locations of these trenches is on sheet 2 (fig. 3).

Data are shown on mosaics of high-resolution color photographs made using a set of techniques and a special device: a camera mounted on a 0.7 m by 1.2 m aluminum frame. The tubular frame can be seen in some places on the photomosaic logs (sheets 1, 2). Digital images from these photos were rectified to the field grid, string lines visible on the logs. Distance (y-axis) of all three trenches is measured from an arbitrary line parallel to the fault, hence the starting point of each trench is some arbitrary non-zero value.

Our previous report focused on evidence of the four youngest earthquakes, which occurred during the past 500 years. In this phase of the investigation, the most useful paleoearthquake information was developed from 500-2500 yr BP materials and structures best exposed in trench 02A (sheet 2). Because these logs are the most important to the study, we have placed the index maps on this sheet. To save on expense, users may chose to print only sheet 2 for some of the most critical results. Sheets 4, 6 and 8 also contain critical information about some paleoearthquakes not shown directly on other logs. See the stratigraphic cross-section (sheet 7) for best overview of the entire stratigraphic section in the pond and the age relationship between earthquake data found on the eastern trace to the paleoearthquakes on the western trace. Sheet 1 (log of 01A) and sheet 3 (logs of pits 02A, B, C) are the least critical to readers seeking an overview

of the earthquake evidence, although they both contain stratigraphic and radiocarbon information essential to the investigation.

We assigned unit numbers to most of the stratigraphic units, from u05 (east trace) and u10 (west trace) at the lowest level exposed in the trenches to u550, the most recent artificial fill. These unit numbers are used to correlate the radiocarbon samples between trench walls (table 1). The pond deposits are generally fine-grained with varying amounts of silt and clay, and some sand, particularly near the main fault trace. The main (western) fault trace is located near meter 5 (sheets 1, 2 and 5). West of the main fault trace are older sand and gravel alluvial deposits of the Niles alluvial cone (California Department of Water Resources, 1967) predating the inception of subsidence at Tyson's Lagoon. We have dated pond deposits near the base of trenches as old as 2500 years BP, but do not yet know the age of inception of the pond. Color is used to highlight some key stratigraphic units on the logs, which are described in Appendix 1.

Earthquakes cause a disruption of the ground surface along the trace of the causative fault. The original ground surface prior to deformation can be shown on a trench log as an paleoearthquake horizon. Paleoearthquake (event) horizons are shown as green dashed lines labeled on the western trace as A1 through A10. The five most recent events (A1-A5) correspond to events E1, E2, E3, E4 and EY respectively as shown in Lienkaemper and others (2002a). Event horizons observed across the eastern trace are referred to as B1, B2 and B3. In the western trenches all three eastern events converge to the same horizon as event A8. Hence, for the entire trench site there are a total of twelve paleoearthquake event horizons in evidence from these and the earlier logs.

## RADIOCARBON DATING

Along with laboratory radiocarbon ages of the samples from the 2001-2003 USGS trenches (table 1), we also include those from William's 1992 trenches W1 (02E herein) and W2).

## ACKNOWLEDGMENTS

This project funded by U.S. Geological Survey, National Earthquake Hazards Reduction Program #7460-09712. Special thanks go to John Rogers of the Alameda County Public Works Agency for permitting access to the site and J.N. Alt for sharing his unpublished information. Reviews by X and Y and improved the clarity of the logs and text.

### Appendix 1. Stratigraphic Unit descriptions

Unit u10, a bluish gray to blue clay is the lowest pond deposit observed in our trenches. It may actually comprise different units that have been subjected to continuous saturation. In trench 02A, blue clay underlies u15, but in the pits 02B and C, the blue clay may be derived by the reduction of units as young as u31. Blue crystals of unknown origin and mineralogy can be seen by 20X hand lens in the freshly exposed blue clay. At the eastern fault trace the basal unit, u05, is a carbonate soil, which may be laterally correlative with u10. Unit u15, is light brown organic silty clay, which becomes progressively more altered with depth to shades of light gray, developing a mottled appearance suggesting the organic materials are being lost by some chemical process, and at greater depth begins a transition toward blue clay. Unit u20 is very fine sandy silt, relatively low in organics except for a nearly continuous charcoal-rich burn layer in the upper 10-20 cm. Unit u31 is brown clay silt, more organic rich near the top. In trench 02A organic-rich layers, u40.1 and 40.3, distinctly overly u31, but in trench 03A the transition is less abrupt, so upper u31 may correlate with u40.1 and u40.3. A key marker unit, u41, the lower shelly layer, has been observed in all trenches at the site, except that near the eastern trace in trench 02E, it has few to no gastropod shells and is instead a 1-m-thick organic-rich clayey silt. Unit 41 on the higher east side of the pond is more organic rich because this side is submerged less and thus less gleyed. Also, the shells here may have been dissolved by percolating groundwater or simply not deposited where ponding occurs for a much shorter time each year. Unit u45 is a gray brown clayey silt with shells, but many fewer gastropods

than in u41 and shells of freshwater clams are also found. The shells in u45 diminish and disappear approaching the main fault on the west; on the east, the unit is pinching out at pit 02D.

Unit u50 is thin (<0.25 m) near the western fault trace because erosion has removed it, but near the east trace consists of a 2-m-thick sequence of sandy silts u50(s/t/u, p, n, l, j, h) that are interbedded with charcoal-rich, organic-rich clayey silts u50(q, o, m, k, i). It is probably also true that the original thickness of u50 was greater adjacent to the east trace, in part because it includes fan deposits that are becoming increasingly distal to the west. Also they are deposited in a synformal structure adjacent to the fault, thus trapping a greater amount of sediment. Unit 61, is a charcoal-rich alluvial sand with stones at its base. The base of u61 is an angular unconformity over most of the pond and is conformable only in trench 02E (W1). Unit u63 is a sandy silt, apparently a flood deposit associated with the stream flow associated with the sandy deposit of u61. U63 is capped by u70, a series of brown clayey silts that in some exposures appear as distinct bands of varying shades of brown and elsewhere seem more massive.

The remaining descriptions refer principally to the section observed in trench 00A (Lienkaemper and others, 2002b), because these units are well-developed there and poorly developed in most subsequent trenches. For convenience, one may refer to the log of trench 01A (sheet 1) in the present report, where these units are fairly well-represented. Unit u80, an orange silty clay contains abundant charcoal and fire ash east of the fault zone. Overlying the orange silty clay are gray silty clays, units u90 and u95, identical except that unit u95 contains several fine charcoal stringers, the lowest of which forms the base of the unit. Above these gray clays is a black, organic silty clay, u100, colored by the abundant charcoal within it. Overlying the black clay is gray clayey silt, u110, which contains none of the gastropods that are abundant in three overlying units. Unit u120 is a silt with many gastropods that varies laterally in color from brown to gray. In trench 00A it overlies a blocky scarp colluvium, unit BSC1. The blocky scarp colluvial units are described in Lienkaemper and others (2002b). Overlying u120 is a similar unit, u130, differing mostly by a greater abundance of gastropods and generally a darker brown color. Both silt units u120 and u130 show considerable lateral variation in relative sand, clay and organic content and bioturbation has obscured much of the contact between them. Unit u135 is a dark brown silt with some gastropods and contains a layer of fire ash and charcoal that is continuous east of the fault zone. Unit u135 is best seen near m 10 on the north wall of trench 00A. The contact between u135 and u130 is mostly obscured by bioturbation on the south wall. Unit u140 is a brown, organic-rich silt that overlies the blocky component of BSC2 and grades into its upper part. Unit u150 is a gray, sandy silt, underlying a markedly different unit, u200, a dark brown organic silty clay. Above u200 is a brown, sandy silt, unit u250, which overlies a blocky scarp colluvium, BSC3, near m 6 in trench 00A. Unit u300 is dark-brown, organic silty clay. Unit u350 is brown sandy silt. Unit u400 is a thin (2-10 cm), dark brown, organic clayey silt. Unit u450 is brown sandy silt. Unit u500 is a brown to dark brown silt that has increasingly greater development of blocky peds eastward from the center of the graben. Recent artificial fills (containing asphalt and a bottle ca. 1960s at its base), unit u550, of 0.5-m to a few meters thick now cover most of the area west of the fault.

## REFERENCES CITED

- California Department of Water Resources, 1967, Evaluation of ground water resources: South Bay (Appendix A: Geology), California Department of Water Resources Bulletin, p. 153.
- Lawson, A.C., 1908, The earthquake of 1868, *in* Lawson, A.C., ed., The California earthquake of April 18, 1906: Report of the State Earthquake Investigation Commission (Volume D): Carnegie Institution of Washington Publication 87, p. 434-448.
- Lienkaemper, J.J., 1992, Map of recently active traces of the Hayward Fault, Alameda and Contra Costa counties, California: U.S. Geological Survey Miscellaneous Field Studies Map MF-2196, 13 p. (1 sheet). [<ftp://clover.wr.usgs.gov/pub/ehz/jlienk/mf2196>]
- Lienkaemper, J.J., Dawson, T.E., Personius, S.F., Seitz, G.G., Reidy, L.M., and Schwartz, D.P., 2002a, Logs and data from trenches across the Hayward Fault at Tyson's Lagoon (Tule Pond), Fremont, Alameda County, California: U.S. Geological Survey Miscellaneous Field Studies Map MF-2386, 12 p. (3 sheets). [<http://geopubs.wr.usgs.gov/map-mf/mf2386/>]

- Lienkaemper, J.J., Dawson, T.E., Personius, S.F., Seitz, G.G., Reidy, L.M., and Schwartz, D.P., 2002b, A record of large earthquakes on the southern Hayward Fault: *Seismological Society of America Bulletin*, v. 92, no. 7, p. 2637-2658.
- Wallace, R.E., 1990, Geomorphic expression, *in* Wallace, R.E., ed., *The San Andreas Fault System, California*: U.S. Geological Survey Professional Paper 1515, p. 14-58.
- Williams, P.L., 1993, Geologic record of southern Hayward Fault earthquakes: *California Division of Mines and Geology Special Publication*, v. 113, p. 171-179.
- Woodward-Clyde and Associates, 1970, Fremont Meadows active fault investigation and evaluation, Fremont, California [on file at California Division of Mines and Geology, San Francisco, California; file no. AP-744].
- Working Group on California Earthquake Probabilities, 1999, Earthquake probabilities in the San Francisco Bay Region: 2000-2030— —A Summary of findings: U.S. Geological Survey Open-File Report 99-517, 55 p. [<http://geopubs.wr.usgs.gov/open-file/of99-517/>]

Table 1  
Radiocarbon ages from charcoal samples

Sample no <sup>1</sup>	<sup>14</sup> C age (yr) BP, corrected <sup>2</sup>	Location			d <sup>13</sup> C <sup>3</sup>	Lab no. <sup>4</sup>
		Unit no.	wall	(m)		
W1-5	1540 ± 30	u50(i)	s	2	-26.42	OS-9102
W1-5a	2780 ± 45	u50(i)	s	2	-26.17	OS-4513
W1-12	2090 ± 30	u50(o)	s	2	-25.63	OS-5140
W1-20	3390 ± 110	u50(j)	s	3	-23.88	OS-9091
W1-32	1910 ± 35	u50(i)	s	4	-26.66	OS-4507
W1-39	1940 ± 35	u50(o)	s	4	-24.41	OS-4506
W1-47	1910 ± 50	u50(r)	s	5	-24.65	OS-4508
W1-58	2520 ± 30	u50(tu)	s	7	-26.83	OS-7543
W1-99	4130 ± 70	u41(v)	s	10	-26.06	OS-9101
W1-100	4120 ± 70	u41(v)	s	10	-25.82	OS-7541
W1-109	2940 ± 55	u50(tu)	s	10	-25.41	OS-7925
W1-113	2690 ± 45	u50(i)	s	1	-25.16	OS-5313
W1-141	6360 ± 140	u05(x)	s	11	-23.33	OS-9096
W1-143	7570 ± 95	u05(x)	n	12	-24.73	OS-9099
W1-154	>Modern	u50(p)	s	2	-25.69	OS-7551
W1-178	2150 ± 25	u50(o)	n	7	-26.1	OS-7542
W1-180	1560 ± 40	u50(i)	n	7	-26.63	OS-7547
W1-183	1760 ± 30	u50(u)	n	8	-24.54	OS-7545
W2-9	3140 ± 45	u50(p)	s	10	-25.66	OS-4514
W2-18	2530 ± 55	u70(d)	s	0	-25.76	OS-4515
W2-22	1310 ± 30	u63(e)	s	2	-25.96	OS-4509
W2-25	3120 ± 75	u50(d)	n	0	-25.62	OS-4516
W2-29	1780 ± 30	u50(q)	s	1	-26.8	OS-4510
W2-30	1910 ± 30	u50(q)	s	1	-26.58	OS-9090
W2-31	2000 ± 30	u50(q)	s	1	-25.87	OS-4511
W2-34	1430 ± 40	u50(p)	n	5	-25.18	OS-4512
W2-58	1810 ± 30	u50(p)	n	2	-25.89	OS-5138
W2-61	1700 ± 40	u63(e)	s	3	-23.85	OS-5139
W2-71	3550 ± 65	u50(q)	s	3	-25.62	OS-5141
W2-71a	3110 ± 50	u50(q)	s	3	-26.52	OS-7535
W2-83	2660 ± 85	u50(r)	s	1	-27.62	OS-9095
01A-02	1510 ± 50	u61	s	8	-24.7	79168
01A-03	1260 ± 40	u61	s	9	-25.9	79169
01A-04	1270 ± 40	u61	s	9	-26.8	79170
01A-05	1300 ± 40	u61	s	9	-26.3	79171
01A-06	1330 ± 40	u61	s	8	-25.9	79172
01A-07	1320 ± 30	u61	s	5	-23.4	79173
01A-08	1350 ± 40	u61	s	5	-28.4	79174
01A-09	1290 ± 40	u61	s	5	-24.7	79175
01A-10	1290 ± 40	u61	n	10	-25.8	79176
03A-1	1970 ± 40	u20	s	8	-25	100715
03A-1a	2740 ± 70	u20	s	8	-26	101210
03A-2	2355 ± 30	u20	s	8	-25.91	100716
03A-2a	2310 ± 35	u20	s	8	-25.23	101211
03A-3	2645 ± 40	u20	s	8	-25	100717
03A-4	3135 ± 35	u20	s	7	-25.89	100718
03A-5	2895 ± 40	u20	s	6	-25	100719
03A-6	2175 ± 45	u31c	s	8	-25	100720
03A-7	1860 ± 35	u31b	s	8	-25.15	100721
03A-8	1970 ± 35	u31b	s	8	-26	100722
03A-9	1965 ± 30	u31b	s	8	-25.23	100723
03A-10	2070 ± 30	u31b	s	9	-25.51	100724
03A-11	3410 ± 45	u31a	s	9	-26.61	100725
03B-1	1610 ± 35	u50(i)	s	2	-25.99	100726

<sup>1</sup>Sample number includes trench designation before the hyphen

<sup>2</sup>Ages corrected for d<sup>13</sup>C (Stuiver and Pollach, 1977)

<sup>3</sup>d<sup>13</sup>C value without decimal are estimated with ±2 (Stuiver and Pollach, 1977)

<sup>4</sup>AMS analyses on early trenches (W1, W2) by National Ocean Sciences AMS Facility (NOAMS), Woods Hole, MA; later ones by Center for Accelerator Mass Spectrometry (CAMS), Lawrence Livermore National Laboratory,